

H.P. BREATH TEST

PATIENT NAME _____

DATE OF TEST _____

PATIENTS SHOULD HAVE NOTHING TO EAT OR DRINK AFTER MIDNIGHT
PATIENTS SHOULD NOT HAVE USED ANTIOTIOTICS FOR 1 MONTH
PATIENTS SHOULD NOT HAVE USED PEPTO-BISMOL FOR 1 MONTH
PATIENTS SHOULD NOT HAVE USED CARAFATE IN THE LAST TWO WEEKS
PATIENT SHOULD BE OFF OF PROTON PUMP INHIBITORS FOR 2 WEEKS

PROTON PUMP INHIBITORS INCLUDE:

1. NEXIUM
2. PRILOSEC
3. PREVACID
4. ACIPHEX
5. PROTONIX

PATIENT CAN USE H-2 BLOCKERS IN THEIR PLACE:

H-2 BLOCKERS INCLUDE:

1. TAGAMET
2. ZANTAC
3. AXID
4. PEPCID

THESE H-2 BLOCKERS CAN BE FOUND OVER THE COUNTER

PATIENTS CAN ALSO USE ANTACIDS AS RESCUE MEDICINES THESE INCLUDE:

1. MAALOX
2. ROLAIDS
3. TUMS
4. MYLANTA
5. GELUSIL